

Computational Seismology: Wave equations

Heiner Igel

Department of Earth and Environmental Sciences
Ludwig-Maximilians-University Munich

Goals of this lecture

- Presenting the basic **wave equations** used in this course
- Demonstrating the differences between **scalar** and **vectorial** wave equations
- Showing **first order** and **second order** wave equations
- Presenting **analytical solutions** for acoustic and elastic wave equations
- Introducing some **fundamental solutions** of elastic waves in homogeneous media

Scalar wave equation: analytical solutions

Scalar wave equation

Acoustic wave equation

$$\partial_t^2 p = c^2 \Delta p + s$$

$p \rightarrow p(\mathbf{x}, t)$, pressure


$c \rightarrow c(\mathbf{x})$, velocity

$s \rightarrow s(\mathbf{x}, t)$, source term

Initial conditions

$$p(\mathbf{x}, t = 0) = p_0(\mathbf{x}, t)$$

$$\partial_t p(\mathbf{x}, t = 0) = 0$$



Snapshot of $p(\mathbf{x}, t)$ (solid line) after some time for initial condition $p_0(\mathbf{x}, t)$ (Gaussian, dashed line), 1D case.

Analytical solution: source-free case

Solution

$$p(x, t) = \frac{1}{2}p_0(ct - x) + \frac{1}{2}p_0(ct + x)$$

The solution shown in the movie has been obtained with a finite-difference approximation.

Acoustic wave equation: external source


Green's Function G

$$\partial_t^2 G(\mathbf{x}, t; \mathbf{x}_0, t_0) - c^2 \Delta G(\mathbf{x}, t; \mathbf{x}_0, t_0) = \delta(\mathbf{x} - \mathbf{x}_0) \delta(t - t_0)$$

Delta function δ

$$\delta(x) = \begin{cases} \infty & x = 0 \\ 0 & x \neq 0 \end{cases}$$

$$\int_{-\infty}^{\infty} \delta(x) dx = 1, \quad \int_{-\infty}^{\infty} f(x) \delta(x) dx = f(0)$$



δ -generating function using boxcars.

Acoustic wave equation: analytical solutions

Green's functions for the inhomogeneous acoustic wave equation for all dimensions. $H(t)$ is the Heaviside function.

1D	2D	3D
$\frac{1}{2c} H\left(t - \frac{ r }{c}\right)$	$\frac{1}{2\pi c^2} \frac{H\left(t - \frac{ r }{c}\right)}{\sqrt{t^2 - \frac{r^2}{c^2}}}$	$\frac{1}{4\pi c^2 r} \delta\left(t - r/c\right)$
$r = x$	$r = \sqrt{x^2 + y^2}$	$r = \sqrt{x^2 + y^2 + z^2}$

Acoustic wave equation: analytical solutions

