


# Orthogonal functions - Function Approximation




- The Problem
- Fourier Series
- Chebyshev Polynomials

---


## The Problem

we are trying to approximate a function  $f(x)$  by another function  $g_n(x)$  which consists of a sum over  $N$  *orthogonal* functions  $\Phi(x)$  weighted by some coefficients  $a_n$ .

$$f(x) \approx g_N(x) = \sum_{i=0}^N a_i \Phi_i(x)$$



# The Problem




... and we are looking for optimal functions in a least squares ( $L_2$ ) sense ...

$$\|f(x) - g_N(x)\|_{L_2} = \left[ \int_a^b \{f(x) - g_N(x)\}^2 dx \right]^{1/2} = \text{Min!}$$


... a good choice for the basis functions  $\Phi(x)$  are *orthogonal* functions.  
*What are orthogonal functions?* Two functions  $f$  and  $g$  are said to be orthogonal in the interval  $[a,b]$  if

$$\int_a^b f(x)g(x)dx = 0$$

How is this related to the more conceivable concept of orthogonal vectors? Let us look at the original definition of integrals:




# Orthogonal Functions - Definition





$$\int_a^b f(x)g(x)dx = \lim_{N \rightarrow \infty} \left( \sum_{i=1}^N f_i(x)g_i(x)\Delta x \right)$$

... where  $x_0=a$  and  $x_N=b$ , and  $x_i-x_{i-1}=\Delta x$  ...

If we interpret  $f(x_i)$  and  $g(x_i)$  as the  $i$ th components of an  $N$  component vector, then this sum corresponds directly to a scalar product of vectors. The vanishing of the scalar product is the condition for *orthogonality* of vectors (or functions).




$$f_i \bullet g_i = \sum_i f_i g_i = 0$$



# Periodic functions

Let us assume we have a piecewise continuous function of the form


$$f(x+2\pi) = f(x)$$




... we want to approximate this function with a linear combination of  $2\pi$  periodic functions:

$1, \cos(x), \sin(x), \cos(2x), \sin(2x), \dots, \cos(nx), \sin(nx)$

$$\Rightarrow f(x) \approx g_N(x) = \frac{1}{2}a_0 + \sum_{k=1}^N \{a_k \cos(kx) + b_k \sin(kx)\}$$



# Orthogonality of Periodic functions




... are these functions orthogonal ?

$$\begin{aligned} \int_{-\pi}^{\pi} \cos(jx) \cos(kx) dx &= \begin{cases} 0 & j \neq k \\ 2\pi & j = k = 0 \\ \pi & j = k > 0 \end{cases} \\ \int_{-\pi}^{\pi} \sin(jx) \sin(kx) dx &= \begin{cases} 0 & j \neq k, j, k > 0 \\ \pi & j = k > 0 \end{cases} \\ \int_{-\pi}^{\pi} \cos(jx) \sin(kx) dx &= 0 \quad j \geq 0, k > 0 \end{aligned}$$


... YES, and these relations are valid for any interval of length  $2\pi$ .  
Now we know that this is an orthogonal basis, but how can we obtain the coefficients for the basis functions?



from minimising  $f(x)-g(x)$



# Fourier coefficients



optimal functions  $g(x)$  are given if



$$\|g_n(x) - f(x)\|_2 = \text{Min !} \quad \text{or} \quad \frac{\partial}{\partial a_k} \left\{ \|g_n(x) - f(x)\|_2 \right\} = 0$$

... with the definition of  $g(x)$  we get ...

$$\frac{\partial}{\partial a_k} \|g_n(x) - f(x)\|_2^2 = \frac{\partial}{\partial a_k} \left[ \int_{-\pi}^{\pi} \left[ \frac{1}{2} a_0 + \sum_{k=1}^N \{a_k \cos(kx) + b_k \sin(kx)\} - f(x) \right]^2 dx \right]$$

leading to (nice exercise)

$$g_N(x) = \frac{1}{2} a_0 + \sum_{k=1}^N \{a_k \cos(kx) + b_k \sin(kx)\} \quad \text{with}$$
$$a_k = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos(kx) dx, \quad k = 0, 1, \dots, N$$
$$b_k = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin(kx) dx, \quad k = 1, 2, \dots, N$$



# Fourier approximation of $|x|$


... Example ...


$$f(x) = |x|, \quad -\pi \leq x \leq \pi$$

leads to the Fourier Serie


$$g(x) = \frac{1}{2}\pi - \frac{4}{\pi} \left\{ \frac{\cos(x)}{1^2} + \frac{\cos(3x)}{3^2} + \frac{\cos(5x)}{5^2} + \dots \right\}$$

.. and for  $n < 4$   $g(x)$  looks like





# Fourier approximation of $x^2$




... another Example ...

$$f(x) = x^2, \quad 0 < x < 2\pi$$


leads to the Fourier Serie

$$g_N(x) = \frac{4\pi^2}{3} + \sum_{k=1}^N \left\{ \frac{4}{k^2} \cos(kx) - \frac{4\pi}{k} \sin(kx) \right\}$$


.. and for  $N < 11$ ,  $g(x)$  looks like







# Fourier - discrete functions



... what happens if we know our function  $f(x)$  only at the points


$$x_i = \frac{2\pi}{N} i$$

it turns out that in this *particular* case the coefficients are given by


$$\begin{aligned} a_k^* &= \frac{2}{N} \sum_{j=1}^N f(x_j) \cos(kx_j), & k &= 0, 1, 2, \dots \\ b_k^* &= \frac{2}{N} \sum_{j=1}^N f(x_j) \sin(kx_j), & k &= 1, 2, 3, \dots \end{aligned}$$

.. the so-defined Fourier polynomial is the unique interpolating function to the function  $f(x_j)$  with  $N=2m$

$$g_m^*(x) = \frac{1}{2} a_0^* + \sum_{k=1}^{m-1} \{a_k^* \cos(kx) + b_k^* \sin(kx)\} + \frac{1}{2} a_m^* \cos(kx)$$




# Fourier - collocation points




... with the important property that ...

$$g_m^*(x_i) = f(x_i)$$


... in our previous examples ...



$f(x)=|x| \Rightarrow f(x)$  - blue ;  $g(x)$  - red;  $x_i$  - '+'




# Fourier series - convergence






$f(x)=x^2 \Rightarrow f(x)$  - blue ;  $g(x)$  - red;  $x_i$  - '+'

$N = 4$



$N = 8$







# Fourier series - convergence


$f(x)=x^2 \Rightarrow f(x)$  - blue ;  $g(x)$  - red;  $x_i$  - '+'

$N = 16$




$N = 32$






# Orthogonal functions - Gibb's phenomenon





$f(x)=x^2 \Rightarrow f(x)$  - blue ;  $g(x)$  - red;  $x_i$  - '+'


$N = 16$




$N = 32$




$N = 128$




$N = 256$



The overshoot for equi-spaced Fourier interpolations is  $\approx 14\%$  of the step height.



# Chebyshev polynomials



We have seen that Fourier series are excellent for interpolating (and differentiating) periodic functions defined on a regularly spaced grid. In many circumstances physical phenomena which are not periodic (in space) and occur in a *limited area*. This quest leads to the use of **Chebyshev polynomials**.


We depart by observing that  $\cos(n\varphi)$  can be expressed by a polynomial in  $\cos(\varphi)$ :

$$\cos(2\varphi) = 2\cos^2\varphi - 1$$


$$\cos(3\varphi) = 4\cos^3\varphi - 3\cos\varphi$$

$$\cos(4\varphi) = 8\cos^4\varphi - 8\cos^2\varphi + 1$$

... which leads us to the definition:



# Chebyshev polynomials - definition



$$\cos(n\varphi) = T_n(\cos(\varphi)) = T_n(x), \quad x = \cos(\varphi), \quad x \in [-1,1], \quad n \in N$$

... for the Chebyshev polynomials  $T_n(x)$ . Note that because of  $x=\cos(\varphi)$  they are defined in the interval  $[-1,1]$  (which - however - can be extended to  $\mathfrak{R}$ ). The first polynomials are

$$T_0(x) = 1$$


$$T_1(x) = x$$

$$T_2(x) = 2x^2 - 1$$


$$T_3(x) = 4x^3 - 3x$$

$$T_4(x) = 8x^4 - 8x^2 + 1 \quad \text{where}$$


$$|T_n(x)| \leq 1 \quad \text{for} \quad x \in [-1,1] \quad \text{and} \quad n \in N_0$$



# Chebyshev polynomials - Graphical




The first ten polynomials look like [0, -1]




The  $n$ -th polynomial has extrema with values 1 or -1 at

$$x_k^{(ext)} = \cos\left(\frac{k\pi}{n}\right), \quad k = 0, 1, 2, 3, \dots, n$$







# Chebyshev collocation points




These extrema are not equidistant (like the Fourier extrema)



$$x_k^{(ext)} = \cos\left(\frac{k\pi}{n}\right), \quad k = 0, 1, 2, 3, \dots, n$$



# Chebyshev polynomials - orthogonality



... are the Chebyshev polynomials orthogonal?


*Chebyshev polynomials are an orthogonal set of functions in the interval  $[-1, 1]$  with respect to the weight function  $1/\sqrt{1-x^2}$  such that*

$$\int_{-1}^1 T_k(x) T_j(x) \frac{dx}{\sqrt{1-x^2}} = \begin{cases} 0 & \text{for } k \neq j \\ \pi / 2 & \text{for } k = j > 0 \\ \pi & \text{for } k = j = 0 \end{cases}, \quad k, j \in N_0$$


... this can be easily verified noting that

$$x = \cos \varphi, \quad dx = -\sin \varphi d\varphi$$

$$T_k(x) = \cos(k\varphi), \quad T_j(x) = \cos(j\varphi)$$



# Chebyshev polynomials - interpolation




... we are now faced with the same problem as with the Fourier series. We want to approximate a function  $f(x)$ , this time not a periodical function but a function which is defined between  $[-1,1]$ .

We are looking for  $g_n(x)$


$$f(x) \approx g_n(x) = \frac{1}{2} c_0 T_0(x) + \sum_{k=1}^n c_k T_k(x)$$

... and we are faced with the problem, how we can determine the coefficients  $c_k$ . Again we obtain this by finding the extremum (minimum)

$$\frac{\partial}{\partial c_k} \left[ \int_{-1}^1 \{g_n(x) - f(x)\}^2 \frac{dx}{\sqrt{1-x^2}} \right] = 0$$



# Chebyshev polynomials - interpolation



... to obtain ...

$$c_k = \frac{2}{\pi} \int_{-1}^1 f(x) T_k(x) \frac{dx}{\sqrt{1-x^2}}, \quad k = 0, 1, 2, \dots, n$$


... surprisingly these coefficients can be calculated with FFT techniques, noting that

$$c_k = \frac{2}{\pi} \int_0^\pi f(\cos \varphi) \cos k\varphi d\varphi, \quad k = 0, 1, 2, \dots, n$$


... and the fact that  $f(\cos \varphi)$  is a  $2\pi$ -periodic function ...

$$c_k = \frac{1}{\pi} \int_{-\pi}^\pi f(\cos \varphi) \cos k\varphi d\varphi, \quad k = 0, 1, 2, \dots, n$$

... which means that the coefficients  $c_k$  are the Fourier coefficients  $a_k$  of the periodic function  $F(\varphi) = f(\cos \varphi)$ !



# Chebyshev - discrete functions



... what happens if we know our function  $f(x)$  only at the points

$$x_i = \cos \frac{\pi}{N} i$$

in this *particular* case the coefficients are given by


$$c_k^* = \frac{2}{N} \sum_{j=1}^N f(\cos \varphi_j) \cos(k \varphi_j), \quad k = 0, 1, 2, \dots, N/2$$

... leading to the polynomial ...


$$g_m^*(x) = \frac{1}{2} c_0^* T_0 + \sum_{k=1}^m c_k^* T_k(x)$$

... with the property

$$g_m^*(x) = f(x) \quad \text{at} \quad x_j = \cos(\pi j/N) \quad j = 0, 1, 2, \dots, N$$




# Chebyshev - collocation points - $|x|$





$f(x)=|x| \Rightarrow f(x)$  - blue ;  $g_n(x)$  - red;  $x_i$  - '+'

8 points




16 points






# Chebyshev - collocation points - $|x|$





$f(x)=|x| \Rightarrow f(x)$  - blue ;  $g_n(x)$  - red;  $x_i$  - '+'

32 points




128 points






# Chebyshev - collocation points - $x^2$




$f(x)=x^2 \Rightarrow f(x)$  - blue ;  $g_n(x)$  - red;  $x_i$  - '+'

8 points




The interpolating function  $g_n(x)$  was shifted by a small amount to be visible at all!


64 points









# Chebyshev vs. Fourier - numerical



Chebyshev




Fourier




$f(x)=x^2 \Rightarrow f(x)$  - blue ;  $g_N(x)$  - red;  $x_i$  - '+'


**This graph speaks for itself ! Gibb's phenomenon with Chebyshev?**




# Chebyshev vs. Fourier - Gibb's



Chebyshev




Fourier




$f(x)=\text{sign}(x-\pi) \Rightarrow f(x)$  - blue ;  $g_N(x)$  - red;  $x_i$  - '+'


**Gibb's phenomenon with Chebyshev? YES!**




# Chebyshev vs. Fourier - Gibb's




## Chebyshev




## Fourier



$f(x) = \text{sign}(x - \pi) \Rightarrow f(x)$  - blue ;  $g_N(x)$  - red;  $x_i$  - '+'



# Fourier vs. Chebyshev



## Fourier

$$x_i = \frac{2\pi}{N} i$$

periodic functions

$$\cos(nx), \sin(nx)$$

$$g_m^*(x) = \frac{1}{2} a_0^* + \sum_{k=1}^{m-1} \{a_k^* \cos(kx) + b_k^* \sin(kx)\} + \frac{1}{2} a_m^* \cos(kx)$$

*collocation points*

*domain*

*basis functions*

*interpolating function*


## Chebyshev

$$x_i = \cos \frac{\pi}{N} i$$


limited area [-1,1]

$$T_n(x) = \cos(n\varphi), \\ x = \cos \varphi$$

$$g_m^*(x) = \frac{1}{2} c_0^* T_0 + \sum_{k=1}^m c_k^* T_k(x)$$



# Fourier vs. Chebyshev (cont'd)



## Fourier

$$a_k^* = \frac{2}{N} \sum_{j=1}^N f(x_j) \cos(kx_j)$$

$$b_k^* = \frac{2}{N} \sum_{j=1}^N f(x_j) \sin(kx_j)$$

- Gibb's phenomenon for discontinuous functions
- Efficient calculation via FFT
- infinite domain through periodicity

*coefficients*

*some properties*

## Chebyshev

$$c_k^* = \frac{2}{N} \sum_{j=1}^N f(\cos \varphi_j) \cos(k\varphi_j)$$

- limited area calculations
- grid densification at boundaries
  - coefficients via FFT
- excellent convergence at boundaries
- Gibb's phenomenon